



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

roosting and feeding places of the Little Blue Heron and American Egret, etc.—J. A. A.

**Nelson on New Birds from Mexico.**—Ten of the thirteen species and subspecies here described<sup>1</sup> were obtained by Mr. Nelson and his assistant Mr. Goldman during their expedition to southwestern Mexico in the winter of 1902-03, mostly in the States of Guerrero and Michoacan. In most cases the new forms are based on good series of specimens, and several of them seem quite strongly differentiated from their nearest known allies.—J. A. A.

**Nelson's 'Revision of the North American Mainland Species of *Myiarchus*.'**<sup>2</sup>—The present paper covers the species of the genus *Myiarchus* occurring north of the Isthmus of Panama, including those of Cozumel Island and the Tres Marias Islands. Nine species are recognized, with ten additional subspecies, of which three of the latter, belonging to the *lawrencei* group, are described as new. In his introductory remarks Mr. Nelson calls attention to the evanescent character of the brighter or more intense colors of the freshly acquired plumage. "This extreme intensity of coloration [of the fresh plumage] quickly passes into a duller condition which continues with but little change through the winter months. In spring the colors gradually fade or become bleached by the sun until in the breeding season the original shades of greenish, olive and gray of the back and the yellow of the under parts are almost lost in the dingy browns and yellows of the frayed plumage." He also calls attention to the wide range of variation in the extent of the dusky pattern of the tail feathers, the non-recognition of which has led to the recording of *M. nuttingi* as a bird of southern Arizona, the supposed Arizona specimens of *nuttingi* proving to be merely females of *M. cinerascens*. Mr. Nelson, however, adds to the United States list *Myiarchus crinitus residuus* Howe, based on Florida specimens, on the ground of a slight average difference in the length of the bill. This separation had previously been made, on exactly the same basis, by Mr. Bangs and rejected by the A. O. U. Committee as too unimportant for recognition in nomenclature.

Mr. Nelson discusses at some length the old case of *Tyrannula mexicana* Kaup vs. *Myiarchus cooperi* Baird, without reaching a positive conclusion, but gives his reasons for believing that *Tyrannula mexicana* = *Tyrannula cinerascens* Lawrence, and that the present *Myiarchus mexi-*

---

<sup>1</sup> Descriptions of New Birds from Southern Mexico. By E. W. Nelson. Proc. Biol. Soc. Washington, Vol. XVI, pp. 151-160, Nov. 30, 1903.

<sup>2</sup> Revision of the North American Mainland Species of *Myiarchus*. By E. W. Nelson. Proc. Biol. Soc. Washington, Vol. XVII, pp. 21-30, March 10, 1904.